

Brief Report: Impact of Antiretroviral Regimen on Pregnancy and Infant Outcomes in Women With HIV/ HBV Coinfection

Flavia Matovu Kiweewa, PhD, Camlin Tierney, PhD, [...], and Debika Bhattacharya, MD

Background:

There are limited data on the impact of antenatal antiretroviral regimens (ARV) on pregnancy and infant outcomes in HIV/HBV coinfection. We compared outcomes among 3 antenatal antiretroviral regimens for pregnant women with HIV/HBV.

Methods:

The PROMISE study enrolled ARV-naïve pregnant women with HIV. Women with HBV were randomized to (no anti-HBV)-zidovudine (ZDV) + intrapartum nevirapine and 1 week of tenofovir disoproxil fumarate and emtricitabine (TDF-FTC); (3TC)-3TC + ZDV + LPV/r; or (FTC-TDF)-FTC + TDF + LPV/r. Pairwise group comparisons were performed with Fisher exact, *t*, or log rank tests. Adverse pregnancy outcome (APO) was a composite of low birth weight, preterm delivery, spontaneous abortion, stillbirth, or congenital anomaly.

Results:

Of 138 women with HIV/HBV, 42, 48, and 48 were analyzed in the no anti-HBV, 3TC, and FTC-TDF arms. Median age was 27 years. APOs trended lower in the no anti-HBV (26%) vs 3TC (38%), and FTC-TDF arms (35%), $P \geq 0.25$). More infant deaths occurred among the FTC-TDF [6 (13%)] vs no anti-HBV [2 (5%)] and 3TC [3 (7%)] arms. There were no differences in time-to-death, HIV-free survival, birth or one-year WHO Z-score length-for-age, and head circumference. Hepatitis B e antigen (HBeAg) was associated with an increased risk of APO, 48% vs 27% (odds ratio 2.79, 95% confidence interval: 1.19 to 6.67, *post hoc*).

Conclusion:

With HBV/HIV coinfection, the risk of an APO was increased with maternal ARV compared with ZDV alone, although the differences were not statistically significant. Maternal HBeAg was associated with a significantly increased risk of APO. Infant mortality was highest with FTC + TDF + LPV/r. Early assessment of HBeAg could assist in identifying high-risk pregnancies for close monitoring.

Key Words: HIV, HBV, antiretroviral therapy, infant outcomes

INTRODUCTION

About 3%–12% of pregnant women are living with HIV and HBV yet data on the impact of antiretroviral (ARV) regimens on pregnancy and infant outcomes in HIV/HBV coinfection are limited.^{1,2} Tenofovir disoproxil fumarate (TDF) is the preferred agent for treating HBV in pregnancy, given its high barrier to resistance, favorable safety profile, and efficacy for HIV and HBV.^{3–5} Previously a cornerstone ARV, lamivudine (3TC) decreases perinatal HBV transmission and is safe in pregnancy.^{6,7} Similarly, emtricitabine (FTC) possesses anti-HBV activity.⁸

The PROMISE study demonstrated that infant mortality was increased with ritonavir-boosted lopinavir (LPV/r)+TDF/FTC compared with antepartum zidovudine (ZDV) alone, but whether such differences also exist in the context of HBV-active regimens in HIV/HBV coinfection is unknown. In this secondary analysis, our objective was to compare the impact of ARV regimen on pregnancy and infant adverse outcomes among people living with HIV and HBV. We further assessed the association of maternal baseline HBeAg status and HBV viral load with pregnancy and infant outcomes.

METHODS

Study Population and Eligibility Criteria

PROMISE 1077BF/1077FF was a multisite randomized open-label trial examining strategies for preventing perinatal HIV transmission and for preserving maternal and infant health, implemented between April 2011 and September 2016. The protocol was approved by relevant ethics committees in the collaborating countries, and reviewed by an independent data and safety monitoring board. All women provided written informed consent for study participation. The study population included women with high CD4⁺ cell counts ($\geq 350/\text{mm}^3$) who had not met

country-specific criteria for initiating ART. Women with elevated ALT ($>2.5 \times$ ULN) were excluded. Additional details are referenced here.⁹ Women who were HBsAg-positive were randomized 1:1:1 to receive zidovudine (ZDV) plus intrapartum single-dose nevirapine followed by 6–14 days of TDF and FTC postpartum (no anti-HBV), 3TC + ZDV + Lopinavir/ritonavir (LPV/r) (3TC), or FTC + TDF + LPV/r (FTC-TDF).

Clinical and safety evaluations occurred at entry, week 2, 4, 8, every 4 weeks until labor and delivery; at labor and delivery; at postpartum week 1, week 6, 14, and Q12 weeks thereafter. Screening HBsAg was performed at local laboratories. Maternal HBV DNA levels were measured at Quest with COBAS AmpliPrep/COBAS TaqMan HBV Test, v2.0 assay (lower limit of quantification (LLQ) 20 IU/mL), and later with the COBAS HBV/Roche 8800 (LLQ 10 IU/mL) assay.

Adverse pregnancy outcome (APO) was defined as a composite of low birth weight (LBW) (<2500 g), preterm delivery (<37 weeks), spontaneous abortion (<20 weeks), stillbirth (≥ 20 weeks), or a congenital anomaly.⁹ Infant outcomes were: mortality (time to), HIV-free survival (time to earlier of HIV infection or death), weight and hemoglobin level (primary focus at birth), and weight and WHO-Z score for length-for-age (primary focus at 1 year) and head circumference (primary focus at 1 year), grade 3 or 4 adverse events, graded for severity per the DAIDS toxicity table 2009.¹⁰

Pairwise group comparisons between regimens and their outcomes were performed with Fisher exact, t , or log rank tests. The primary comparison was between FTC-TDF and 3TC arms. Infant time-to-event analyses were censored at the earlier of last clinic visit, last laboratory measurement, or end of infant follow-up of visit week 104, or site notification of the START study results, which recommended initiation of ARVs for all women.¹¹

We applied logistic and Cox proportional hazards regression models to further assess *post hoc* whether there was treatment effect modification by maternal baseline HBeAg status and HBV DNA level ($<$ or ≥ 1 million IU/mL), and their association, with APO, infant hemoglobin at birth, birth weight, infant weight at 1 year, and time-to-infant mortality. The statistical significance level was set at 0.05.

RESULTS

Socio-Demographic and Clinical Characteristics

Of 3543 women, 139 (3.9%)¹² were HBsAg-positive and 42, 48, and 49 randomized to the no anti-HBV, 3TC, and FTC-TDF-containing arms, Table 1. One participant was excluded from analyses. Half of the women had a $CD4 \geq 500$ cells/mm³ (51%). Median baseline HBV DNA level was 2.6 log₁₀ IU/mL; 23% had HBV DNA ≥ 1 million IU/mL, whereas 24% had a level < 20 IU/mL. Thirty-four (26%) women were HBeAg-positive, and among these, 28 (93%) of 30 with data had HBV VL ≥ 1 million IU/mL (Table 1).

| Characteristic | No anti-HBV (N = 42) | 3TC (N = 48) | FTC |
|----------------|----------------------|--------------|-----|
| Country | | | |
| South Africa | 9 (21%) | 10 (20%) | 1 |
| Malawi | 19 (45%) | 20 (42%) | 1 |
| Zambia | 1 (2%) | 2 (4%) | 0 |
| Uganda | 4 (10%) | 4 (8%) | 0 |
| Zimbabwe | 7 (17%) | 6 (13%) | 0 |

TABLE 1.

Baseline Characteristics of Pregnant Women Living With HIV and HBV in the PROMISE Study

Pregnancy Outcomes

Among 132 (96%) of 138 pregnancies with outcomes, there were no spontaneous abortions, 5 singleton stillbirths, and 127 pregnancies with 126 live singleton births and 1 twin birth (in FTC-TDF arm). The risk of an APO was not statistically different in the primary pairwise comparison of the FTC-TDF vs 3TC arms (35% and (38%) of pregnancies, estimated odds ratio {[OR, 3TC reference] 0.86, 95% confidence interval (CI): [0.34 to 2.17], $P = 0.83$, Table 2}, but were numerically higher when compared with no anti-HBV arm (26%); FTC-TDF vs no anti-HBV [1.55 (0.55 to 4.47), $P = 0.48$; 3TC vs no anti-HBV 1.80 (0.65 to 5.13), $P = 0.25$]. The most common APOs were preterm delivery and LBW. Among 127 live birth pregnancies, preterm delivery (<37 weeks' gestation) was 11% in the no anti-HBV arm and higher in 3TC and FTC-TDF arms with 20% and 28% [OR (95% CI) 2.12 (0.52 to 10.26) and 3.25 (0.87 to 14.92)] The proportion with LBW (<2500 g) was 16% (6/37) in the no anti-HBV arm and, 20% (9/44) and 17% (8/46) in the 3TC and FTC-TDF arms, and did not differ between arms, $P \geq 0.78$.

| Variable | Total (N = 132) | No anti-HBV (N = 42) | 3TC (N = 47) | FTC-TDF (N = 43) | P |
|-----------------|-----------------|----------------------|--------------|------------------|---|
| No. of outcomes | | | | | |
| Singleton | 126 | 30 (71%) | 47 | 49 (95%) | |
| Twin | 1 | 0 | 1 | 0 | |
| Total | 127 | 30 (24%) | 48 | 49 (39%) | |

TABLE 2.

Pregnancy Outcomes Among Women Living With HIV and HBV in the PROMISE Study

Infant Outcomes

Through 2 years, more infant deaths occurred among the FTC-TDF [6 (13%)] vs no anti-HBV [2 (5%)] and 3TC [3 (7%)] arms. More infant deaths occurred among the FTC-TDF [6 (13%)] vs no anti-HBV [2 (5%)] and 3TC [3 (7%)] arms, Figure 1, Supplemental Digital Content, <http://links.lww.com/QAI/B914>. Two, 1, and 0 deaths in the FTC-TDF, 3TC, and no anti-HBV arms were preterm. One death in each of the 3TC and FTC-TDF arms occurred in infants born at 34 weeks of gestation (very preterm delivery infants). One, 2, and 0 deaths in each arm had LBW (<2500 g), and none had very LBW (<1500 g). There were no significant differences in time-to-death, HIV-free survival, or mean WHO Z-score length-for-age, and head circumference at birth or 1 year.

The risk of a grade 3 or 4 infant adverse event was not different between arms. Infants in the FTC-TDF arm had significantly higher mean hemoglobin level at birth vs those in 3TC (1.31 g/dL, 95% CI: 0.38 to 2.25, $P < 0.01$), but not vs no anti-HBV arm (-0.77 g/dL, 95% CI: -1.75 to 0.22 , $P = 0.12$); 3TC vs no anti-HBV arm (0.54 g/dL, 95% CI: -0.39 to 1.48 , $P = 0.25$, see Figure 2 and Tables 1 and 2, Supplemental Digital Content, <http://links.lww.com/QAI/B914>.

Pregnancy and Infant Outcomes by Maternal HBeAg Status and Baseline HBV Viral Load

In *post hoc* analyses, treatment effects did not differ by maternal HBeAg for APO, infant hemoglobin level at birth, birth weight, weight at 1 year, and time-to-infant mortality ($P \geq 0.12$, see Table 1, Supplemental Digital Content, <http://links.lww.com/QAI/B914>). Adjusted for treatment arm, HBeAg positivity was associated with an increased risk of an APO [48% (15/31) vs 27% (25/94)] in HBeAg-negative women [OR 2.79 (95% CI: (1.19 to 6.67), $P = 0.019$)], Table 3, Supplemental Digital Content, <http://links.lww.com/QAI/B914>.

Higher percentages were observed for pregnancies among HBeAg-positive women for each of stillbirth (10% vs 2%), preterm (32% vs 15%) and LBW (21% vs 14%). Results were similar when we compared outcomes by baseline maternal HBV DNA above and below 1 million IU/mL, Tables 2 and 3, Supplemental Digital Content, <http://links.lww.com/QAI/B914>. There was no evidence of an association between HBeAg or maternal HBV DNA and infant hemoglobin at birth, infant birth weight, and infant weight at 1 year, Table 3, Supplemental Digital Content, <http://links.lww.com/QAI/B914>.

DISCUSSION

This is one of the first studies to evaluate whether ARV regimens affected pregnancy and infant outcomes in HIV/HBV coinfection. The proportion of women with any APO was numerically higher with antepartum TDF containing ART, although differences were not statistically significant. Infant mortality at 2 years was higher in the TDF-ART arm, with two-thirds of deaths occurring within the first month of life in ART arms. However, event numbers were small. Although these results were not statistically significant, they follow previous results from similar results from the overall PROMISE trial. Use of 3TC plus ZDV resulted in significantly lower infant mean hemoglobin at birth compared with TDF-ART. Maternal baseline HBeAg positivity and HBV DNA ≥ 1 million IU/mL were associated with increased risk of an APO.

Similar point estimates for APO treatment differences were observed in the much larger parent PROMISE study, where higher rates of APO were observed with 3TC + ZDV + LPV/r and FTC + TDF + LPV/r than with ZDV alone, but not between the 2 PI-containing ART regimens.

Although the exact mechanisms responsible for the relationship between maternal ART and the elevated risk of APO are unclear, one possible mechanism is through modulation of the immune response. Use of ART may induce APOs by altering circulating cytokine levels through ART-associated immune reconstitution.^{13,14} In addition, use of protease inhibitors during pregnancy is associated with decreased progesterone levels, suggesting a potential mechanism contributing to LBW and preterm deliveries¹⁵ as observed in the PROMISE primary analysis.⁹

Similar to the PROMISE study, we observed that infant mortality was highest with FTC + TDF + LPV/r initiated during pregnancy. In HIV/HBV coinfection, one may have hypothesized that non-HBV active therapy would have led to worse infant outcomes given 2 viral infections in pregnancy, but we did not see this effect, and instead, saw the same interesting association with infant mortality with TDF-ART. As in HIV, the mechanism is unknown but may be related to an additive deleterious impact of TDF and LPV/r. Of note, we did not assess the impact of postpartum maternal ART status on infant outcomes, and factors beyond antepartum ARV receipt may have contributed to infant outcomes.

In *post hoc* analyses, APOs were more pronounced among women with positive HBeAg, consistent with other studies.^{16,17} Although the mechanism is unclear, maternal HBeAg is associated with high HBV viral load.¹⁸ HBeAg serves as a qualitative and cheaper surrogate marker of viral load.¹⁹ Data from HPTN-046, a randomized controlled of HIV vertical transmission in sub-Saharan Africa that evaluated 6 months of infant nevirapine vs. placebo for perinatal HIV prevention showed that HBsAg-positive women with high HBV viral loads (defined as HBV VL ≥ 1

million IU/mL) had a higher risk of delivering a LBW infant compared with those with HBV VL < 1 million IU/mL. Similarly, in a large retrospective cohort study of 26,350 pregnant Thai women, 1446 (5.5%) of whom had HBV, preterm births and LBW were higher among women with positive HBeAg status (21.9% vs 14.1%, $P = 0.005$) and (8.6% vs 13.6%, $P = 0.028$).¹⁷

Major strengths of our study include the randomized design, including a no anti-HBV ZDV-alone regimen, and the relatively long two-year infant follow-up. We also include data from a variety of SSA countries where WHO has noted a paucity of data.³ Study limitations include *post hoc* analyses, inability to distinguish between acute and chronic HBV infection with a single timepoint assessment, and the use of LPV/r as backbone for the regimen containing TDF + 3TC. This agent is being replaced by dolutegravir as a first-line regimen, and therefore extrapolating our results to other TDF + 3TC-containing regimens needs caution. Finally, this analysis had a relatively small sample size and few events, limiting our precision on estimating treatment effects.

In HIV/HBV coinfection, maternal ART (3TC + ZDV + LPV/r and FTC + TDF + LPV/r) initiated during pregnancy was associated with higher risk of APO compared to use of ZDV alone though the difference were not statistically significant.⁹ Baseline positive HBeAg or high baseline HBV DNA (≥ 1 million IU/mL) was associated with higher risk for an APO. Early antenatal diagnosis of HBV, followed by assessment of HBeAg in pregnancy as a surrogate for high HBV DNA levels, particularly in resource limited settings, could assist in identifying high-risk pregnancies for close monitoring. As was demonstrated in PROMISE overall, among this subgroup of women with HIV and HBV, infant mortality was also highest with FTC + TDF + LPV/r. Future research should evaluate the safety of newer ART/HBV drug-containing regimens including tenofovir alafenamide fumarate and dolutegravir in pregnant women living with HIV/HBV.²⁰

Supplementary Material

[Click here to view.](#)^(251K, docx)

ACKNOWLEDGMENTS

The authors thank the study participants, funders, and the PROMISE Study staff who implemented the study.

The authors thank the mothers and their infants who participated, the PROMISE staff at the clinical research sites; the staff at the IMPAACT central laboratory at the University of North Carolina at Chapel Hill; the statistical staff at the Center for Biostatistics in AIDS Research, Harvard T.H. Chan School of Public Health; and the data-management staff at the Frontier Science and Technology Research Foundation.

The study products were provided free of charge by Abbott, Gilead Sciences, Boehringer Ingelheim, and GlaxoSmithKline.

Footnotes

D.B. and F.M. report research grant support from Gilead Sciences Inc, under award numbers IN-US-987-5950 and GPHA 07844 respectively). M.G.P. has been a consultant to Aligos and Antios. Overall support for the International Maternal Pediatric Adolescent AIDS Clinical Trials Network (IMPAACT) was provided by the National Institute of Allergy and Infectious Diseases (NIAID) with co-funding from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) and the National Institute of Mental Health (NIMH), all components of the National Institutes of Health (NIH), under Award Numbers UM1AI068632 (IMPAACT LOC), UM1AI068616 (IMPAACT SDMC) and UM1AI106716 (IMPAACT LC), and by NICHD contract number HHSN275201800001I. Additional funding from NIAID UM1AI069465 to A.G., NIAID R01 01AI100748-01 and NICHD R01 HD085862 to D.B. Study products were provided free by AbbVie, Gilead Sciences, Boehringer Ingelheim, and ViiV/GlaxoSmithKline. Award Numbers UM1AI068632-15 (IMPAACT LOC), UM1AI068616-15 (IMPAACT SDMC) and UM1AI106716-15 (IMPAACT LC), and by NICHD contract number HHSN275201800001I. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. The other authors have no conflicts of interest to disclose.

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site (www.jaids.com).

Article information

J Acquir Immune Defic Syndr. 2022 Sep 1; 91(1): 79–84.

Published online 2022 May 27. doi: [10.1097/QAI.0000000000003022](https://doi.org/10.1097/QAI.0000000000003022)

PMCID: PMC9377493

NIHMSID: NIHMS1806899

PMID: 35621877

Flavia Matovu Kiweewa, PhD,^{a,b} Camlin Tierney, PhD,^c Kevin Butler, MS,^c Marion G. Peters, MD,^d Tichaona Vhembo, MD,^e Dhayendre Moodley, PhD,^f Vani Govender, BSc,^g Neaka Mohtashemi, BA,^h Hannah Ship, BA,ⁱ Philippa Musoke, MD,^{aj} Dingase Dula, MBBS,^k Kathy George, MPH,^l Nahida Chakhtoura, MD,^m Mary G. Fowler, MD,ⁿ Judith S. Currier, MD,^h and Debika Bhattacharya, MD^h

^aMakerere University—Johns Hopkins University Research Collaboration, Kampala, Uganda;

^bDepartment of Epidemiology and Biostatistics, Makerere University School of Public Health, College of Health Sciences, Kampala, Uganda;

^cCenter for Biostatistics in AIDS Research, Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA;

^dDepartment of Medicine, Feinberg School of Medicine, Northwestern University CRS, Chicago, IL;

^eDepartment of Obstetrics and Gynaecology, University of Zimbabwe College of Health Sciences, Harare, Zimbabwe;

^fDepartment of Obstetrics and Gynaecology, University of KwaZulu-Natal, Nelson Mandela School of Medicine, Durban, South Africa;

^gCentre for the AIDS Programme of Research in South Africa, Congella, South Africa;

^hDepartment of Medicine, Division of Infectious Diseases, David Geffen School of Medicine, University of California, Los Angeles, CA;

ⁱUniversity of Miami Miller School of Medicine, Miami, FL;

^jDepartment of Paediatrics and Child Health, College of Health Sciences, Makerere University;

^kCentre for AIDS Research in South Africa and Department of Obstetrics and Gynecology, School of Clinical Medicine, University of KwaZulu Natal, Durban, South Africa;

^lFamily Health International 360, Durham, NC;

^mNational Institutes of Health (NIH), Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD), Bethesda, MD; and

ⁿDepartment of Pathology, Johns Hopkins University School of Medicine, Baltimore, MD

 Corresponding author.

Camlin Tierney: tierney@sdac.harvard.edu; Kevin Butler: kbutler@sdac.harvard.edu; Marion G. Peters: Marion.Peters@ucsf.edu; Tichaona Vhembo:

tvhembo@uz-ctrc.org; Dhayendre Moodley: Moodleyd1@ukzn.ac.za; Vani Govender: Vani.Govender@caprisa.org; Neaka Mohtashemi: NMohtashemi@mednet.ucla.edu;

Hannah Ship: hannah.ship@med.miami.edu; Philippa Musoke: pmusoke@mujhu.org; Dingase Dula: ddula@jhu.medcol.mw; Kathy George: KGeorge@fhi360.org; Nahida

Chakhtoura: nahida.chakhtoura@nih.gov; Mary G. Fowler: mfowler5@jhmi.edu; Judith S. Currier: JSCurrier@mednet.ucla.edu; Debika Bhattacharya:

debikab@mednet.ucla.edu

Correspondence to: Flavia Matovu Kiweewa, PhD, Department of Epidemiology and Biostatistics, Makerere University—Johns Hopkins University Research

Collaboration, Makerere University College of Health Sciences, School of Public Health, Old Mulago Hill Road, Kampala, Uganda (e-mail: fmatovu@musph.ac.ug).

Received 2021 Sep 2; Accepted 2022 Apr 28.

Copyright © 2022 The Author(s). Published by Wolters Kluwer Health, Inc.

This is an open access article distributed under the terms of the [Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 \(CCBY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

REFERENCES

1. Bayo P, Ochola E, Oleo C, et al.. High prevalence of hepatitis B virus infection among pregnant women attending antenatal care: a cross-sectional study in two hospitals in northern Uganda. *BMJ Open*. 2014;4(11):e005889. [PMC free article] [PubMed] [Google Scholar]
2. Liu J, Zhang S, Liu M, et al.. Maternal pre-pregnancy infection with hepatitis B virus and the risk of preterm birth: a population-based cohort study. *Lancet Glob Health*. 2017;5:e624–e632. [PubMed] [Google Scholar]
3. World Health Organization. Prevention of Mother-To-Child Transmission of Hepatitis B Virus: Guidelines on Antiviral Prophylaxis in Pregnancy: Web Annex A: Systematic Review of the Efficacy and Safety of Antiviral Therapy during Pregnancy. 2020. [PubMed] [Google Scholar]
4. Pan CQ, Mi LJ, Bunchorntavakul C, et al.. Tenofovir disoproxil fumarate for prevention of vertical transmission of hepatitis B virus infection by highly viremic pregnant women: a case series. *Dig Dis Sci*. 2012;57:2423–2429. [PMC free article] [PubMed] [Google Scholar]
5. Liver EAFTSOT. EASL 2017 Clinical Practice Guidelines on the management of hepatitis B virus infection. *J Hepatol*. 2017;67:370–398. [PubMed] [Google Scholar]
6. Funk AL, Lu Y, Yoshida K, et al.. Efficacy and safety of antiviral prophylaxis during pregnancy to prevent mother-to-child transmission of hepatitis B virus: a systematic review and meta-analysis. *Lancet Infect Dis*. 2021;21:70–84. [PubMed] [Google Scholar]

7. Han L, Zhang HW, Xie JX, et al.. A meta-analysis of lamivudine for interruption of mother-to-child transmission of hepatitis B virus. *World J Gastroenterol.* 2011;17:4321. [PMC free article] [PubMed] [Google Scholar]
8. Matthews GV, Seaberg E, Dore GJ, et al.. Combination HBV therapy is linked to greater HBV DNA suppression in a cohort of lamivudine-experienced HIV/HBV co-infected individuals. *AIDS.* 2009;23:1707. [PMC free article] [PubMed] [Google Scholar]
9. Fowler MG, Qin M, Fiscus SA, et al.. Benefits and risks of antiretroviral therapy for perinatal HIV prevention. *N Engl J Med.* 2016;375:1726–1737. [PMC free article] [PubMed] [Google Scholar]
10. U.S. Department of Health and Human Services, National Institutes of Health, National Institute of Allergy and Infectious Diseases, Division of AIDS. Division of AIDS Table for Grading the Severity of Adult and Pediatric Adverse Events, Version 1.0. [Updated August 2009]. Available at: <https://rsc.niaid.nih.gov/sites/default/files/table-for-grading-severity-of-adult-pediatric-adverse-events.pdf> [Google Scholar]
11. Lundgren JD, Babiker AG, Gordin F, et al.. Initiation of antiretroviral therapy in early asymptomatic HIV infection. *N Engl J Med.* 2015;373:795–807. [PMC free article] [PubMed] [Google Scholar]
12. Bhattacharya D, Gupta A, Tierney C, et al. Hepatotoxicity and liver-related mortality in women of childbearing potential living with human immunodeficiency virus and high CD4 cell counts initiating Efavirenz-containing regimens. *Clin Infect Dis.* 2021;72(8):1342–1349. [PMC free article] [PubMed] [Google Scholar]
13. Fiore S, Newell M-L, Trabattoni D, et al.. Antiretroviral therapy-associated modulation of Th1 and Th2 immune responses in HIV-infected pregnant women. *J Reprod Immunol.* 2006;70:143–150. [PubMed] [Google Scholar]
14. Fiore S, Ferrazzi E, Newell M-L, et al.. Protease inhibitor—associated increased risk of preterm delivery is an immunological complication of therapy. *J Infect Dis.* 2007;195:914–916. [PubMed] [Google Scholar]
15. Papp E, Mohammadi H, Loutfy MR, et al.. HIV protease inhibitor use during pregnancy is associated with decreased progesterone levels, suggesting a potential mechanism contributing to fetal growth restriction. *J Infect Dis.* 2015;211:10–18. [PMC free article] [PubMed] [Google Scholar]
16. Bhattacharya D, Guo R, Tseng C-H, et al.. Maternal HBV viremia and association with adverse infant outcomes in women living with HIV and HBV. *Pediatr Infect Dis J.* 2021;40:e56–e61. [PMC free article] [PubMed] [Google Scholar]
17. Sirilert S, Traisrisilp K, Sirivatanapa P, et al.. Pregnancy outcomes among chronic carriers of hepatitis B virus. *Int J Gynecol Obstet.* 2014;126:106–110. [PubMed] [Google Scholar]
18. Lu Y, Song Y, Zhai X, et al.. Maternal Hepatitis B e Antigen can be an Indicator for Antiviral Prophylaxis of Perinatal Transmission of Hepatitis B Virus. *Emerg Microbes Infect.* 2021;1–28. [PMC free article] [PubMed] [Google Scholar]
19. Sun KX, Li J, Zhu FC, et al.. A predictive value of quantitative HBsAg for serum HBV DNA level among HBeAg-positive pregnant women. *Vaccine.* 2012;30:5335–5340. [PubMed] [Google Scholar]
20. Gish RG, Elmofti Y. Tenofovir Alafenamide Fumarate—A New Silver Bullet to Prevent Mother-to-child Transmission of Hepatitis B Virus: Wiley Online Library; 2020. [PubMed] [Google Scholar]